

## Contents

Introduction .....	2
Kit Contents .....	3
Before Starting .....	3
Protocol 1: E.Z.N.A.® Plasmid Midi Kit Spin Protocol .....	4
■ Growth of bacterial culture .....	4
■ Lyse bacterial cells with alkaline-SDS Solution .....	4
■ Purify Plasmid with HiBind™ DNA Midi Column .....	5
■ Elution Plasmid DNA From Column .....	6
■ Alternative Elution Step From Column .....	7
Protocol 2: E.Z.N.A.® Plasmid Midi Kit Vacuum Protocol .....	8
Protocol 3: Low Copy-Number Plasmid Protocol .....	9
Yield and Quality of DNA .....	9
Plasmid Copy-Number and Expected Yield .....	10
Trouble Shooting Guide .....	11
Order Information .....	12

Revised October 2005

## Introduction

The E.Z.N.A.® family of products is an innovative system that radically simplifies extraction and purification of nucleic acids from a variety of sources. Key to the system is Omega Bio-Tek's proprietary HiBind® matrix that avidly, but reversibly, binds DNA or RNA under certain optimal conditions, allowing proteins and other contaminants to be removed. Nucleic acids are easily eluted with deionized water or low salt buffer.

The E.Z.N.A.® Plasmid Midi Kit combines the power of HiBind® technology with the time-tested consistency of alkaline-SDS lysis of bacterial cells to deliver high quality plasmid DNA. Omega Bio-Tek's midi-columns facilitate the binding, washing, and elution steps, thus enabling multiple samples to be simultaneously processed.

Yields vary according to plasmid copy number, *E.coli* strain, and conditions of growth, but 50 ml of overnight culture in LB medium typically produces 100-250µg high-copy number plasmid or 10-100 µg Low-copy number plasmid. Up to 100 ml culture may be processed when working with low-copy number plasmids. The product is suitable for automated fluorescent DNA sequencing, restriction endonuclease digestion, transfection of mammalian cells, and other manipulations.

## Storage and Stability

All E.Z.N.A.® Plasmid isolation components are guaranteed for at least 12 months from the date of purchase when stored as follows: Solution I/RNase A at 4°C, all other material at 22-25°C.

## Kit Contents

Product Number	D6904-00	D6904-01	D6904-03	D6904-04
Purification times	2 Preps	10 Preps	25 Preps	100 Preps
HiBind® DNA Midi Column	2	10	25	100
15 ml collection tube	2	10	25	100
Solution I	6 ml	30 ml	70 ml	250 ml
Solution II	6 ml	30 ml	70 ml	250 ml
Solution III	8 ml	40 ml	100 ml	2 x 180 ml
Buffer GPS	5 ml	25 ml	60 ml	250 ml
Buffer HB	8 ml	40 ml	85 ml	2 x 170 ml
DNA Wash Buffer	4 ml	20 ml	50 ml	4 x 50 ml
RNase A	pre-added	150 µl	350 µl	1.2 ml
Elution Buffer	8 ml	20 ml	60 ml	2 x 120 ml
Instruction Booklet	1	1	1	1

## Before Starting

Briefly examine this booklet and become familiar with each step. Prepare all components and have the necessary materials ready before starting.

**Supplied By User:** High speed centrifuge capable of 12,000 × g  
 Laboratory centrifuge equipped with **swinging-bucket** rotor capable of 3,000-5,000 × g.  
 Nuclease-free 15 ml centrifuge tubes  
 High speed centrifuge tubes (polycarbonate or Corex®)  
 95-100% ethanol

<b>IMPORTANT</b>	1. Add vial of RNase A to bottle of Solution I provided and store at 4°C.
	2. <b>DNA Wash Buffer</b> is to be diluted with 96-100% ethanol as follows:
	D6904-00 Add 16 ml 96-100% ethanol
	D6904-01 Add 80 ml 96-100% ethanol
	D6904-03 Add 200 ml 96-100% ethanol
	D6904-04 Add 200 ml 96-100% ethanol per bottle
<b>Store diluted DNA Wash Buffer at room temperature</b>	

## Protocol 1: E.Z.N.A.® Plasmid Midi Kit Spin Protocol

This Protocol is designed to isolate 100-250 µg of high Copy-Number plasmids or 10-100 µg of low Copy-Number Plasmids from 30-50 ml overnight cultures. For increasing yield of low Copy-Number plasmid, proceed as “Low Copy-Number Plasmids protocol” on page 8.

### ■ Growth of bacterial culture

1. **Culture volume: Inoculate 30-50 ml LB/ampicillin (50 µg/ml) medium placed in a 0.2-0.4 liter culture flask with *E. coli* carrying desired plasmid and grow at 37°C with agitation for 12-16 hours.** For best results use overnight culture as the inoculum. It is strongly recommended that an *endA* negative strain of *E. coli* be used for routine plasmid isolation. Examples of such strains include DH5α® and JM109®.

Optimal growth conditions of bacteria is vital of obtaining maximal plasmid DNA yields. The best conditions are achieved by picking a single isolated colony from a freshly transformed or freshly plate to inoculate a 2-5ml starter culture containing the appropriate antibiotic. Incubate for ~8 hours at 37°C with vigorous shaking (~300 rpm). Then used to inoculate appropriate volume of Pre-warmed liquid growth medium with antibiotic. Grow at 37°C for 12-16 hours with vigorous shaking(~300rpm).Using a flask or vessel with a volume of a least 3-4 times the volume of the culture and dilute the starter culture 1/500 to 1/1000 into growth medium.

Following overnight bacterial growth, an OD<sub>600</sub> of 1.5~2.0 indicates a well-grown culture. For the best result determination of OD<sub>600</sub> for each culture is recommended. it is important to dilute the bacterial culture (10 to 20 fold) to enable photometric measurement in the linear range between 0.1 and 0.5 OD<sub>600</sub>. For maximal yields, the OD<sub>600</sub> of cultures should be under 3.0.

If using a frozen glycerol stock as inoculum, streak it onto an agar plate containing the appropriate antibiotic for single colony isolation. Then picking a single colony and inoculate the 2-5ml starter culture as described above.

- **Lyse bacterial cells with alkaline-SDS Solution**
2. **Pellet up to 30-50 ml bacteria in appropriate vessels by centrifugation at 3,500-5,000 x g for 10 min at room temperature.**
3. Decant or aspirate medium and discard. To ensure that all traces of the medium are removed, use a clean paper towel to blot excess liquid from the wall of the vessel.

To the bacterial pellet add 2.5 ml Solution I/RNase A. Resuspend cells completely by vortexing or pipetting up and down. *Complete resuspension of cell pellet is vital for obtaining good yield.*

4. Transfer cell suspension to a 15 or 50 ml centrifuge tube capable of withstanding 12,000 × g (screw-cap polycarbonate or Corex® glass tubes will suffice). **Add 2.5 ml Solution II, cover, and mix gently but thoroughly by inverting and rotating tube 10-15 times to obtain a cleared lysate.** A 2 min incubation at room temperature may be necessary.

Avoid vigorous mixing as this will shear chromosomal DNA and lower plasmid purity. Prolonged incubation may lead to nicking of plasmid DNA. (Store Solution II tightly capped when not in use.)

5. **Add 3.2 ml Solution III, cover, and gently mix by inverting tube several times until a flocculent white precipitate forms.** Centrifuge at  $\geq 12,000 \times g$  for 10 min at room temperature (preferably at 4°C) to pellet the cellular debris and genomic DNA.

The Buffers must be mixed thoroughly. If the mixture appears still viscous, brownish and conglobated, more mixing is required to completely neutralize the solution. Complete neutralization of the solution is vital of obtaining good yields. Increasing centrifugation speed is helpful to completely remove the precipitated bacterial cell material. A tightly packed cell debris pellet indicates efficient lysis.

Note: For fast removing the precipitated bacterial cell material, one may order OBI Lysate Clearance filter syringe to replace centrifugation of this step. This filter cartridge completely removes SDS precipitates and clears bacterial lysates using filter syringe. Besides, one may order the whole kit, E.Z.N.A. Fastfiler Plasmid midi or maxi kits.

***Step 6 to 13 should be performed in swinging-bucket rotor for maximal plasmid DNA yields. And all centrifugation steps must be carried out at room temperature.***

- **Purify Plasmid DNA with HiBind™ DNA Midi Column**
- 6. **Prepare the HiBind Midi Column.** Place a HiBind Midicolumn into a 15 ml collection tube, provided. Add 2 ml of Buffer GPS to the column and let it sit at room temperature for 3-10 min. Spin in a swinging bucket rotor at 3,000-5,000 xg for 5 min at room temperature. Discard the eluate and assemble the column in the 15 ml collection tube.

7. **Carefully aspirate and add 3.7 ml of the clear supernatant to the HiBind® DNA Midi column assembled in the 15 ml collection tube, making sure that no cellular debris is carried over.** The Midi column has a maximum capacity of 4 ml. Centrifuge at 3,000-5,000 × g for 3-5 min at room temperature to completely pass lysate through column. Discard the flow-through liquid and repeat this step until the entire sample has been passed through. Finally discard the flow-through and reuse the collection tube in Step 8.

8. **Add 3 ml Buffer HB to the Midi column and centrifuge as above.** This step ensures that residual protein contamination is removed and must be included for downstream applications requiring high quality DNA. Discard flow-through liquid and reuse the collection tube in the next step.

9. **Wash the column by adding 3.5 ml of DNA Wash Buffer diluted with ethanol.** Centrifuge as above and discard flow-through.

Note: DNA Wash Buffer Concentrate must be diluted with absolute ethanol before use. See label for directions. If refrigerated, DNA Wash Buffer must be brought to room temperature before use.

10. **Optional Step: Repeat wash step with another 3.5 ml DNA Wash Buffer.** Centrifuge as above and discard fluid.

11. **Centrifuge the empty capped column for 10-15 min at maxi speed (no more than 8,000 x g) to dry the column matrix.**

*DO NOT skip this step - it is critical for removing traces of ethanol that may otherwise interfere with downstream applications.*

- **Elution Plasmid DNA From HiBind™ DNA Midi column.**  
**Optional:** For maximal yield and high concentration of plasmid, see alternative protocol of elution on page 7. For fast elution, proceed step 12-13.

12. **Further Drying The Column (Optional).** Choose either of the methods below to further dry the column before eluting DNA (only if necessary):

- A. Place the column into a vacuum container to dry the ethanol for 10 min. Then, remove the column and place into a vacuum chamber at room temperature. Any device connected to a vacuum source may be used. Seal the chamber and apply vacuum for 15 min. Remove the column and proceed to Step 13.

B. Bake the column in a vacuum oven or incubator at 65°C for 10 min. Remove the column and proceed to Step 13.

13. **Place column into a clean 15 ml centrifuge tube. Add 0.5-1.0 ml (depending on desired concentration of final product) Elution Buffer (or TE buffer) directly onto the column matrix.** Allow column to sit 2 min at room temperature. Centrifuge at max speed (no more than 8,000 x g) for 5 min to elute DNA. This represents approximately 60-80% of bound DNA. An optional second elution will yield any residual DNA, though at a lower concentration. Alternatively, a second elution may be performed using the first eluate to maintain a high DNA concentration. Also, preheating the water to 70°C prior to elution may significantly increase yields.

Note: The plasmid DNA obtained using this protocol performs well in PCR, restriction digests, lipid mediated transfection and transformation. The expected concentration of plasmid is vary between different copy number vector. However, the concentration of high copy-number plasmid is 150-400 ug/ml. Some residual ethanol may present, but does not interfere with these downstream applications. One may get high concentration and absolutely remove ethanol with optional elution step as following.

■ **Alternative protocol of Elution Plasmid from Column**

1. **Place HiBind™ DNA Midi column into a clean 15 ml centrifuge tube. Add 3 ml Elution Buffer (or TE buffer) directly onto the column matrix.** Allow column to sit 2 min at room temperature. Centrifuge for 5 min at max speed to elute DNA.
2. **Carefully transfer eluted plasmid from 15 ml centrifuge tube to a clean tube suitable for precipitation. add 130ul 5M NaCl and 2.2 ml room temperature isopropanol. Vortex to mix and centrifuge at >15,000 x g for 30 min at 4°C.** Carefully decant the supernatant.
3. **Wash DNA pellet once one with 1ml ice-cold 70% ethanol and centrifuge at > 15,000 x g for 10 min.** Carefully decant the supernatant without disturbing the pellet and air-dry the pellet for 10-15 min.
4. **Finally resuspend DNA pellet in 200-500µl (depending on desired concentration of final product) Elution Buffer or TE Buffer.**

## Protocol 2. E.Z.N.A.® Plasmid Midi Kit Vacuum Protocol

1. **Prepare cleared cell lysate by following step 1-5 of centrifugation procedure on page 4-5.**
2. **Prepare the vacuum manifold according to manufacturer's instructions and connect the Midi column to the manifold.**
3. Add **2 ml** of Buffer GPS to the column and let it sit at room temperature for 3-10 min. Apply the vacuum to allow all the liquid pass through the column.
4. **Transfer cleared cell lysate into the HiBind® DNA Midi column**, be careful not to overfill the column, apply the vacuum to allow all sample pass through the column. Repeat transfer the lysate into the column until the entire sample has been passed through.
5. **Add 3.0 ml HB buffer to the column** and apply the vacuum to draw the liquid through the column.
6. **Wash the column: add 3.5 ml of DNA wash buffer (pre-diluted with absolute ethanol) into the column** and allow it pass through the column.
7. **Wash the column again with 3.5 ml DNA wash buffer by repeating step 6.**
8. **Keep the vacuum on for another 10-15 min after the liquid pass through the column.** (This step to ensure the removal of residue ethanol).
9. **Proceed Elution Step as Centrifugation protocol Fast Elution Step on Page 6 or Alternative elution step on page 7.**

### Protocol 3: Low Copy-Number Plasmids Protocol

Low copy plasmids generally give 0.1-1 µg DNA per ml overnight culture. For isolation of plasmid from low copy-number plasmids (0.1-1 µg/ml culture) or low-midi copy-number plasmids (1-2 µg/ml culture) bacteria, the method can be modified to essentially increase the yield if necessary.

Start with 50-100 ml bacterial culture, centrifuge for 10 min at 3,500-5,000 xg in a centrifuge tube. Proceed to Step 3 (Page 4) and double the volumes of Solution I, II, and Solution III. Continue as above using only one HiBind® DNA Midi column. There is no need to increase the volumes of Buffer HB and DNA Wash Buffer used. The Buffer of Solution I, II, III can be purchase separately.

Note: This method is not recommended for high copy number plasmids because above 50 ml culture, the HiBind® DNA Midi column quickly becomes saturated. In this situation we recommend processing of multiple samples from the same culture.

### Yield and quality of DNA

Determine the absorbance of an appropriate dilution (10- to 20-fold) of the sample at 260 nm and then at 280 nm. The DNA concentration is calculated as follows:

$$DNA\ concentration = Absorbance_{260} \times 50 \times (Dilution\ Factor) \mu g/ml$$

The ratio of  $(Absorbance_{260})/(Absorbance_{280})$  is an indication of nucleic acid purity. A value greater than 1.8 indicates > 90% nucleic acid. Alternatively, quantity (as well as quality) can sometimes best be determined by agarose gel/ethidium bromide electrophoresis by comparison to DNA samples of known concentrations. Typically, the majority of the DNA eluted is in monomeric supercoil form, though concatamers may also be present.

### Plasmid Copy-Number and Expected Yield

The Yield and quality of the plasmid DNA depend on a number of factors including plasmid copy number, size of insert, host strain, culture volume, culture medium and binding capacity of kit. In these factors, The copy number of vector, culture volume and binding capacity of kit are most important. Copy number of plasmid is vary from one copy to several hundred copies per cell as dicated by their origin of replication. But very large plasmids are often maintained at very low copy number per cell.

Plasmid	Replicon	Copy Number	expected Yield of 50ml culture
pUC vector	pMB1	500-700	180-220 µg
pBR322 and its derivatives	pMB1	15-20	10-20µg
PACYC and its derivatives	p15A	10-12	5-30 µg
pSC101 and its derivatives	pSC101	~5	5-20 µg
pBluescript	ColE14	300-500	100-150 µg
ColE14	ColE14	15-20	5-20 µg
pGEM	pMB1	300-700	150-200 µg

## Trouble Shooting Guide

Problem	Likely Cause	Suggestions
Low DNA yields	Poor cell lysis	<p>Only use LB or YT medium containing ampicillin. Do not use more than 50 ml with high copy plasmids.</p> <p>Cells may not be dispersed adequately prior to addition of Solution II. Vortex cell suspension to completely disperse.</p> <p>Increase incubation time with Solution II to obtain a clear lysate.</p> <p>Solution II if not tightly closed, may need to be replaced. Prepare as follows: 0.2 M NaOH, 1% SDS.</p>
	Bacterial culture overgrown or not fresh.	Do not incubate cultures for more than 16 hours at 37°C. Storage of cultures for extended periods prior to plasmid isolation is detrimental.
	Low copy-number plasmid used	Such plasmids may yield as little as 0.1µg DNA from a 1 ml overnight culture. Increase culture volume to 100 ml.
No DNA eluted.	DNA Wash Buffer Concentrate not diluted with absolute ethanol.	Prepare Wash Buffer Concentrate as instructed on the label.
High molecular weight DNA contamination of product.	Over mixing of cell lysate upon addition of Solution II.	Do not vortex or mix aggressively after adding Solution II. Adequate mixing is obtained by simply inverting and rotating tube to cover walls with viscous lysate.
Optical densities do not agree with DNA yield on agarose gel.	Trace contaminants eluted from column increase $A_{260}$ .	Make sure to wash column as instructed. Alternatively, rely on agarose gel/ethidium bromide electrophoresis for quantitation.
RNA visible on agarose gel.	RNase A not added to Solution I.	Add 1 vial of RNase to each bottle of Solution I.
Plasmid DNA floats out of well while loading agarose gel	Ethanol not completely removed from column following wash steps.	Centrifuge column as instructed in Step 10 to dry.
Plasmid DNA will not perform in downstream application	Traces of ethanol remain on column prior to elution.	The column must be centrifuged as instructed in Step 10 and dried before elution. Ethanol precipitation may be required following elution

## Ordering Information

Product	Applications	Cat. No.
<b>Standard E.Z.N.A.™ Plasmid Isolation System</b>		
Plasmid Mini Kit I	Isolation of up to 30 µg Plasmid DNA	D6942/3
Plasmid Mini Kit II	Isolation of up to 70 µg Plasmid DNA	D6945
Plasmid Midi Kit	Isolation of up to 250 µg Plasmid DNA	D6904
Plasmid Maxip Kit	Isolation of up to 1.5 mg Plasmid DNA	D6922
Fastfilter Plasmid Midi Kit	Isolation of up to 250 µg Plasmid DNA, featuring filter syringes for lysate clearance	D6905
Fastfilter Plasmid Maxi Kit	Isolation of up to 1.5 mg Plasmid DNA, featuring filter syringes for lysate clearance	D6924
E-Z 96 Fastfilter Plasmid Kit	Isolation of Plasmid DNA using a 96-well format	D1097
E-Z 96 SE Plasmid Kit	Isolation of plasmid DNA using a single plate	D1095
Yeast Plasmid Isolation Kit	Isolation fo Yeast Plasmid DNA	D3476
<b>E.Z.N.A.™ Endotoxin Free Plasmid Isolation System</b>		
Endo-Free Plasmid Mini Kit I	Isolation of up to 30 µg Endotoxin free Plasmid	D6948
Endo-Free Plasmid Mini Kit II	Isolation of up to 70 µg Endotoxin free Plasmid	D6950
Endo-Free Plasmid Mid Kit	Isolation of up to 250 µg Endotoxin free Plasmid DNA, featuring filter syringes for lysate clearance	D6915
EndoFree Plasmid Maxi Kit	Isolation of up to 1.5 mg Endotoxin free Plasmid DNA, featuring filter syringes for lysate clearance	D6926
<b>E.Z.N.A.™ H P Plasmid Isolation System</b>		
HP Plasmid Mini Kit I	Isolation of up to 30 µg of High Purity Plasmid	D7042
HP Plasmid MidiKit	Isolation of up to 200 µg of High Purity Plasmid	D7004
HP Plasmid Maxi Kit	Isolation of up to 1.5 mg of High Purity Plasmid	D7022
<b>E.Z.N.A.™ Single Strand Phage DNA Isolation Kits</b>		
M13 Isolation Kit	Isolation of up to 15 µg of single stranded phage	D6900
E-Z 96 M13 Isolation Kit	Isolation of up to 15 µg of M-13 DNA using a 96-	D1900
<b>E.Z.N.A.™ Large Construct DNA Isolation Kits</b>		
BAC/PAC DNA Isolation Kit	Effective purification of BAC or PAC DNA	D2156
BAC/PAC DNA Isolation Kit	purification of BAC or PAC using a 96-well format	D1056